

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/336590753>

# Whole-exome sequencing in Russian children with non-type 1 diabetes mellitus reveals a wide spectrum of genetic variants in MODY-related and unrelated genes

Article in *Molecular Medicine Reports* · October 2019

DOI: 10.3892/mmr.2019.10751

CITATIONS

0

READS

88

30 authors, including:



**Oleg Glotov**

Saint Petersburg State University

54 PUBLICATIONS 145 CITATIONS

[SEE PROFILE](#)



**Elena A Serebryakova**

Saint Petersburg State University

11 PUBLICATIONS 16 CITATIONS

[SEE PROFILE](#)



**Olga A Efimova**

D. O. Ott Research Institute of Obstetrics, Gynecology and Reproductology

25 PUBLICATIONS 157 CITATIONS

[SEE PROFILE](#)



**Andrey Glotov**

D. O. Ott Research Institute of Obstetrics and Gynecology

87 PUBLICATIONS 399 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Genome Russia [View project](#)



Prognostic Markers in Patients with Colorectal Cancer [View project](#)

# Whole-exome sequencing in Russian children with non-type 1 diabetes mellitus reveals a wide spectrum of genetic variants in MODY-related and unrelated genes

OLEG S. GLOTOV<sup>1-3\*</sup>, ELENA A. SEREBRYAKOVA<sup>1-3\*</sup>, MARIIA E. TURKUNOVA<sup>4\*</sup>, OLGA A. EFIMOVA<sup>1</sup>, ANDREY S. GLOTOV<sup>1-3</sup>, YURY A. BARBITOFF<sup>2,5</sup>, YULIA A. NASYKHOVA<sup>1,2</sup>, ALEXANDER V. PREDEUS<sup>2,5</sup>, DMITRII E. POLEV<sup>2</sup>, MIKHAIL A. FEDYAKOV<sup>2,3</sup>, IRINA V. POLYAKOVA<sup>2,3</sup>, TATYANA E. IVASHCHENKO<sup>1</sup>, NATALIA Y. SHVED<sup>1-3</sup>, ELENA S. SHABANOVA<sup>1</sup>, ALENA V. TISELKO<sup>1</sup>, OLGA V. ROMANOVA<sup>3</sup>, ANDREY M. SARANA<sup>2,3</sup>, ANNA A. PENDINA<sup>1</sup>, SERGEY G. SCHERBAK<sup>2,3</sup>, EKATERINA V. MUSINA<sup>1</sup>, ANASTASIIA V. PETROVSKAIA-KAMINSKAIA<sup>2</sup>, LIUBOV R. LONISHIN<sup>5,6</sup>, LILIYA V. DITKOVSKAYA<sup>4</sup>, LIUDMILA A. ZHELENINA<sup>4</sup>, LUDMILA V. TYRTOVA<sup>4</sup>, OLGA S. BERESENEVA<sup>4,7</sup>, ROSTISLAV K. SKITCHENKO<sup>5,8</sup>, EVGENII N. SUSPITSIN<sup>4</sup>, ELENA B. BASHNINA<sup>7</sup> and VLADISLAV S. BARANOV<sup>1,2</sup>

<sup>1</sup>D.O. Ott Research Institute of Obstetrics, Gynecology and Reproductology; <sup>2</sup>St. Petersburg State University, 199034 St. Petersburg; <sup>3</sup>City Hospital Number 40, Sestroretsk, 197706 St. Petersburg; <sup>4</sup>St. Petersburg State Pediatric Medical University, 194100 St. Petersburg; <sup>5</sup>Bioinformatics Institute, 197342 St. Petersburg; <sup>6</sup>Peter The Great St. Petersburg Polytechnic University, 195251 St. Petersburg; <sup>7</sup>North-Western State Medical University Named After I.I. Mechnikov, 191015 St. Petersburg; <sup>8</sup>ITMO University, Kronverkskiy Prospekt, 49, 197101 St. Petersburg, Russia

Received March 16, 2019; Accepted August 28, 2019

DOI: 10.3892/mmr.2019.10751

**Abstract.** The present study reports on the frequency and the spectrum of genetic variants causative of monogenic diabetes in Russian children with non-type 1 diabetes mellitus. The present study included 60 unrelated Russian children with non-type 1 diabetes mellitus diagnosed before the age of 18 years. Genetic variants were screened using whole-exome sequencing (WES) in a panel of 35 genes causative of maturity onset diabetes of the young (MODY) and transient or permanent neonatal diabetes. Verification of the WES results was performed using PCR-direct sequencing. A total of 38 genetic variants were identified in 33 out of 60 patients (55%). The majority of patients (27/33, 81.8%) had variants in MODY-related genes: *GCK* (n=19), *HNF1A* (n=2), *PAX4* (n=1), *ABCC8* (n=1), *KCNJ11* (n=1), *GCK+HNF1A* (n=1), *GCK+BLK* (n=1) and *GCK+BLK+WFS1* (n=1). A total of 6 patients (6/33, 18.2%) had variants in MODY-unrelated genes: *GATA6* (n=1),

*WFS1* (n=3), *EIF2AK3* (n=1) and *SLC19A2* (n=1). A total of 15 out of 38 variants were novel, including *GCK*, *HNF1A*, *BLK*, *WFS1*, *EIF2AK3* and *SLC19A2*. To summarize, the present study demonstrates a high frequency and a wide spectrum of genetic variants causative of monogenic diabetes in Russian children with non-type 1 diabetes mellitus. The spectrum includes previously known and novel variants in MODY-related and unrelated genes, with multiple variants in a number of patients. The prevalence of *GCK* variants indicates that diagnostics of monogenic diabetes in Russian children may begin with testing for MODY2. However, the remaining variants are present at low frequencies in 9 different genes, altogether amounting to ~50% of the cases and highlighting the efficiency of using WES in non-*GCK*-MODY cases.

## Introduction

Monogenic diabetes accounts for 1-6% of pediatric diabetes patients with the highest incidence among patients manifesting non-type 1 diabetes mellitus in childhood or adolescence (1).

A large, clinically heterogeneous group of dominantly inherited disorders linked to primary  $\beta$ -cell dysfunction is classified as maturity onset diabetes in the young (MODY). To date, 13 genes causative of 13 types of MODY are known (2). MODY is typically diagnosed before 25 years of age; it is non-insulin dependent and its symptoms are usually mild. However, due to the variety of clinical forms caused by a wide spectrum of mutations in MODY-related genes, different treatment strategies are used: From appropriate diet and physical activity to oral and/or insulin therapy.

*Correspondence to:* Dr Oleg S. Glotov, D.O. Ott Research Institute of Obstetrics, Gynecology and Reproductology, 3 Mendeleevskaya Line, 199034 St. Petersburg, Russia  
E-mail: olglotov@mail.ru

\*Contributed equally

**Key words:** monogenic diabetes, maturity onset diabetes of the young, whole-exome sequencing, genetic variants, children

Monogenic diabetes also includes a number of non-MODY transient or permanent neonatal forms occurring under 6 months of age. More than 20 genes are known to be related to congenital neonatal diabetes (3). Depending on the gene involved, neonatal diabetes may follow patterns of dominant or recessive inheritance and may be isolated or associated with a variety of syndromic features (4). However, due to a very early onset of diabetes, hyperglycemia is often diagnosed prior to other syndromic features. The treatment strategy for non-MODY neonatal diabetes depends on the specific genetic defect causing the diabetic phenotype.

Molecular genetic testing is highly recommended for patients suspected of monogenic diabetes as it allows tailoring treatments to specific etiological mechanisms. Up until recently, search for diabetes-related mutations was usually performed by Sanger sequencing and was therefore limited to only a few genes, leaving a considerable proportion of cases without a known cause. Moreover, a number of studies have demonstrated that frequencies of certain monogenic diabetes subtypes vary strongly among different populations (5), challenging the development of unified recommendations for the target gene choice. An efficient technology to detect previously known and to reveal novel mutations related to monogenic diabetes is next-generation sequencing. This technique allows for a rapid analysis of an unlimited number of genes and may provide valuable knowledge on the genetic variants causative of monogenic diabetes in different populations.

Here, using targeted whole-exome sequencing (WES), we studied the frequency and the spectrum of genetic variants causative of monogenic diabetes in a cohort of Russian children with non-type 1 diabetes mellitus.

## Materials and methods

**Study group.** A total of 60 unrelated patients with diabetes and impaired glucose tolerance (pre-diabetes) were prospectively included in the study. All the patients were of Russian ethnicity and resided in Northwest Russia. In accordance with the guidelines of the American Diabetes Association (6), the diagnoses were based on plasma glucose criteria, either the fasting plasma glucose (FPG) and/or the 2-h plasma glucose (2-h PG) value after a 75-g oral glucose tolerance test (OGTT) and/or the HbA1C criteria. All the patients had an onset of diabetes before the age of 18 years and a detectable C-peptide secretion (or a detectable insulin level in the absence of insulin therapy) and were negative for insulin-, islet-cell-, tyrosinphosphatase IA2-, and glutamate decarboxylase-autoantibodies. The exclusion criterion was the presence of the already confirmed syndrome associated with impaired glucose metabolism (such as Prader-Willi syndrome). In 59 cases, family history was available, and in 41 of them, it was positive for diabetes. All the patients were referred to the study by their medical supervisors.

**Sample preparation and whole-exome sequencing.** Genomic DNA was extracted from whole blood by Magna Pure System (Roche) using the standard protocol. Exome DNA libraries were prepared from 100 ng DNA using TruSeq® Exome Sample Preparation kit (Illumina, Inc.), following

the manufacturer's instructions. Libraries were sequenced on Illumina HiSeq 2500 in 2x100 PE mode. An average of 63.6 million sequencing reads were obtained for each sample, yielding ~50x mean coverage of CDS regions and an average of 89% of CDS bases covered at least 10x.

**Bioinformatic analysis.** Bioinformatic analysis of the WES data was done using a pipeline based on bwa v.0.7.12-r1044 aligner, Picard tools v.2.0.1, and Genome Analysis Tool kit (GATK) v.3.5 software with all the necessary preprocessing steps required by the GATK Best Practices workflow (<https://software.broadinstitute.org/gatk/best-practices/>) (7,8). Target enrichment metrics were collected using the Picard CalculateHsMetrics tool. Variant calling was done using GATK HaplotypeCaller in the cohort genotyping mode with 250 samples included into the cohort (samples with a similar ethnical background from St. Petersburg State University Biobank were used for cohort padding). Variants were filtered using variant quality score recalibration (VQSR) and annotated with SnpEff and SnpSift tools (version 4.2). Additional annotations included the following information: rsID of known variants from dbSNP (build 146), allele frequency (AF) from large sequencing consortia-1000 Genomes (9), Exome Aggregation Consortium (ExAC) (10), and ESP6500 (11); and pathogenicity predictions by Polyphen-2 (12), SIFT (13), PROVEAN (14) obtained from dbNSFP database (15) and by Human Splicing Finder (16) and DDIG (17). For additional prediction of protein stability changes caused by missense mutations with uncertain significance, I-Mutant 2.0 (18) was used. Variant ranking was done using a custom scoring metric. Reference minor allele presence in target genes was analyzed using RMA Hunter (19).

To check the possible presence of copy-number variants (CNVs), we analyzed the sequencing coverage across all targeted exons of interest. To this end, we calculated coverage for each interval using GATK, and then normalized the coverage matrix across samples and intervals. We then used z-score value of the normalized coverage to assess the statistical significance of the results.

**Verification of the WES results and family analysis.** Verification of the WES results in probands and subsequent family analyses were performed by PCR-direct sequencing. Specific primers were designed for verification of each case. The PCR products were purified with 5M NH<sub>4</sub>Ac and 96% ethanol and then with 70% ethanol, dried at 60°C, and dissolved in 10 µl of deionized water. After purification, the PCR products were sequenced using an ABI PRISM BigDyeTerminator 3.1 kit reagent (Applied Biosystems). Then, a capillary electrophoresis was performed in a GA3130x1 Genetic Analyzer (Applied Biosystems). Sequences were analyzed using the Sequence Scanner software (Applied Biosystems).

**Analysis of the GCK promoter for c.-71G>C genetic variant.** A single-base substitution c.-71G>C in the GCK promoter is known to be linked to MODY2 phenotype (20). However, WES did not allow for analysis of the GCK promoter for c.-71G>C. For this reason, the GCK promoter was analyzed for c.-71G>C genetic variant by PCR-direct sequencing as described above with the use of Hae III endonuclease and the

following primers: F-5'-GCATGGCAGCTCTAATGACAG G-3' and R-5'-CATCCTAGCCTGCTTCCCTGG-3'.

## Results

*Genetic variants causative of monogenic diabetes in Russian children with non-type 1 diabetes mellitus.* Using whole-exome sequencing followed by PCR-direct sequencing, we identified the frequency and the spectrum of genetic variants causative of monogenic diabetes in 60 Russian children with non-type 1 diabetes mellitus. Genetic variants were screened for a total of 35 genes: 13 genes causative of MODY [*HNF4A* (MODY1), *GCK* (MODY2), *HNF1A* (MODY3), *PDX1* (MODY4), *HNF1B* (MODY5), *NEUROD1* (MODY6), *KLF11* (MODY7), *CEL* (MODY8), *PAX4* (MODY9), *INS* (MODY10), *BLK* (MODY11), *ABCC8* (MODY12), and *KCNJ11* (MODY13)] and 22 genes causative of transient or permanent neonatal diabetes, including the ones related to specific syndromes (*EIF2AK3*, *RFX6*, *WFS1*, *ZFP57*, *FOXP3*, *AKT2*, *PPARG*, *APPL1*, *PTF1A*, *GATA4*, *GATA6*, *GLIS3*, *IER3IP1*, *LMNA*, *NEUROG3*, *PAX6*, *PLAGL1*, *SLC19A2*, *SLC2A2*, *SH2B1*, *SERPINB4*, and *MADD*).

Overall, 33 out of 60 patients (55%) had genetic variants in the target genes (Table I; 21-40). For 12 patients, parents were available for genetic testing and origins of genetic variants were determined. In 11 cases, genetic variants had been inherited from the parents, and in one case, a *de novo* genetic variant was confirmed. Of 33 patients, 27 (81.8%) had genetic variants in MODY-related genes. The majority of these patients (19 out of 27) had genetic variants in *GCK* (MODY2). The spectrum of *GCK* genetic variants included 13 missense mutations, 3 nonsense mutations, 1 in-frame and 3 frameshift deletions, and 1 single-base substitution in the promoter. In two *GCK* mutation-positive patients, two genetic variants were present: Missense mutation along with a single-base substitution in the promoter (patient #27) and missense mutation along with a nonsense mutation (patient #78). The spectrum of the identified *GCK* genetic variants is shown in Fig. 1. Missense mutations in *HNF1A* (MODY3) were registered in two patients. The other MODY-related genetic variants included three cases of missense mutations: In *PAX4* (MODY 9), in *ABCC8* (MODY12), and in *KCNJ11* (MODY 13).

The presence of genetic variants in different target genes was detected in three patients. In one of them, a *GCK* in-frame deletion was accompanied by an *HNF1A* missense mutation (patient #226). In another one, two missense mutations were present: In *GCK* and in *BLK* (patient #529). In the third patient (#662), a splicing defect in *GCK* and missense mutations in *BLK* and *WFS1* were present.

Genetic variants causative of non-MODY monogenic diabetes were found in 6 out of 33 mutation-positive patients (18.2%). These included a nonsense mutation in *GATA6*, three cases of missense mutations in *WFS1*, one case of a homozygous *EIF2AK3* nonsense mutation (patient #411), and one case of missense mutation and a frameshift deletion present in *SLC19A2* (c.164delC and c.161C>A) (patient #432). The *EIF2AK3* nonsense mutations had been inherited from consanguineous parents who were heterozygous carriers of the same mutation. The *SLC19A2* mutations also appeared to have been inherited from the parents: C.164delC from the mother

and c.161C>A from the father, indicating that both *SLC19A2* alleles in patient #432 were affected.

Considering that monogenic diabetes may be associated with deletions and duplications, we analyzed the possible presence of CNVs in the target genes. We found no evidence for CNVs in the target genes in either sample. However, it should be noted that the limitations of WES technology do not allow for confident detection of small-scale CNVs.

*Relationship between genetic variants and diabetic phenotypes.* We analyzed the relationship of the detected genetic variants to the patients' diabetic phenotypes. Among the 38 detected genetic variants, 23 had been previously reported as linked to monogenic diabetes and 15 were novel ones (Table I). According to the American College of Medical Genetics and Genomics (ACMG) guidelines (41), most of the detected genetic variants (18 previously reported and 6 novel ones) were classified as pathogenic or likely pathogenic and thus were considered as causative of the diabetic phenotypes in the studied patients. However, the relationship of the detected *KCNJ11* missense mutation to the diabetic phenotype was not apparent, because earlier it had been shown to be associated with hyperinsulinism (35), which was not present in patient #134.

Three previously reported and 9 novel genetic variants were classified as those of uncertain significance, and two genetic variants were likely benign (Table I). These variants included 12 missense mutations; for them, we performed an additional *in silico* analysis using I-Mutant 2.0 (18) (Table II). In all but one case, the *in silico* modeling attested to a decrease of protein stability, thus suggesting the pathogenic effect of the checked genetic variants. Of special interest were two novel *WFS1* genetic variants, initially classified as likely benign. Patient #266 inherited the genetic variant from a non-diabetic mother, while patient #408 inherited the genetic variant from a mother with diabetes. Homozygous mutations in *WFS1* lead to the development of Wolfram syndrome, an autosomal recessive disorder characterized by a list of clinical signs including a bilateral progressive optic atrophy, deafness, and diabetes mellitus (42). Heterozygous carriers of *WFS1* mutations have been reported to have risk of early-onset diabetes mellitus (43). The latter cannot be excluded in our patients. However, an intriguing point is that the *WFS1* genetic variant in patient #408, who inherited it from a diabetic mother, appeared to not decrease the protein stability according to I-Mutant, which makes its pathogenicity questionable.

*Clinical picture in patients with multiple genetic variants.* Finally, we analyzed the clinical picture in patients with more than one genetic variant in one or different target genes (Table III). A simultaneous presence of two *GCK* genetic variants in patient #27 raised the question of their location in one or both alleles. The parents were not available for analysis. The clinical picture was mild and typical for MODY2. It contrasted with the severe one usually reported in patients with both *GCK* alleles affected (44,45), suggesting that, in patient #27, both genetic variants were present in the same allele and thus had no accumulative effect. In patient #78, who was also a carrier of two *GCK* genetic variants, the clinical picture was typical for MODY2. As both genetic variants were inherited from

Table I. Genetic variants identified in Russian children with non-type 1 diabetes mellitus.

Patient number	Gene	Nucleotide change (protein change)	Mutation type	Mutation origin	Pathogenicity according to ACMG	(Refs.)
59	<i>GCK</i>	c.772G>T (p.Gly258Cys)	Missense	Unknown	Likely pathogenic	(21)
62	<i>GCK</i>	c.930_931delGG (p.Asp311fs)	Frameshift	Unknown	Pathogenic	(22)
83	<i>GCK</i>	c.930_931delGG (p.Asp311fs)	Frameshift	Unknown	Pathogenic	(22)
95	<i>GCK</i>	c.130G>A (p.Gly44Ser)	Missense	Father	Likely pathogenic	(23)
167	<i>GCK</i>	c.128G>A (p.Arg43His)	Missense	Mother	Likely pathogenic	(24)
197	<i>GCK</i>	c.233T>C (p.Leu77Pro)	Missense	Father	Likely pathogenic	(25)
426	<i>GCK</i>	c.683C>T (p.Thr228Met)	Missense	Unknown	Likely pathogenic	(26)
460	<i>GCK</i>	c.682A>G (p.Thr228Ala)	Missense	Mother	Likely pathogenic	(21)
580	<i>GCK</i>	c.775G>A (p.Ala259Thr)	Missense	Unknown	Likely pathogenic	(27)
663	<i>GCK</i>	c.1079C>A (p.Ser360*)	Nonsense	Unknown	Pathogenic	(28)
665	<i>GCK</i>	c.660C>A (p.Cys220*)	Nonsense	Unknown	Pathogenic	(24)
176	<i>GCK</i>	c.1349C>T (p.Ala450Val)	Missense	Unknown	Likely pathogenic	(29)
661	<i>GCK</i>	c.1349C>T (p.Ala450Val)	Missense	Unknown	Likely pathogenic	(29)
118	<i>GCK</i>	c.117_119delAAG (p.Lys39del)	In-frame deletion	Unknown	Uncertain significance	Novel
119	<i>GCK</i>	c.1346_1347delCG (p.Ala449fs)	Frameshift	Unknown	Pathogenic	Novel
434	<i>GCK</i>	c.868G>C (p.Glu290Gln)	Missense	Mother	Uncertain significance	Novel
578	<i>GCK</i>	c.1253G>C (p.Ser418Thr)	Missense	Unknown	Pathogenic	Novel
27	<i>GCK</i>	c.754T>C (p.Cys252Arg)	Missense	Unknown	Likely pathogenic	(30)
		c.-71G>C	Promoter	Unknown	Likely pathogenic	(20)
78	<i>GCK</i>	c.199G>T (p.Glu67*)	Nonsense	Mother	Pathogenic	Novel
		c.766G>C (p.Glu256Lys)	Missense	Mother	Likely pathogenic	(31)
153	<i>HNF1A</i>	c.709A>G (p.Asn237Asp)	Missense	Unknown	Uncertain significance	(32)
422	<i>HNF1A</i>	c.485T>G (p.Leu162Arg)	Missense	Unknown	Uncertain significance	Novel
215	<i>PAX4</i>	c.574C>A (p.Arg192Ser)	Missense	Unknown	Uncertain significance	(33)
114	<i>ABCC8</i>	c.4139G>A (p.Arg1380His)	Missense	Unknown	Likely pathogenic	(34)
134	<i>KCNJ11</i>	c.406C>A (p.Arg136Ser)	Missense	Unknown	Uncertain significance	(35)
68	<i>GATA6</i>	c.1477C>T (p.Arg493*)	Nonsense	De novo	Pathogenic	(36)
266	<i>WFS1</i>	c.2452C>T (p.Arg818Cys)	Missense	Mother	Likely benign	(37)
408	<i>WFS1</i>	c.2327A>T (p.Glu776Val)	Missense	Mother	Likely benign	(38)
133	<i>WFS1</i>	c.1124G>A (p.Arg375His)	Missense	Unknown	Uncertain significance	Novel
411	<i>EIF2AK3</i>	c.1912C>T (p.Arg638*)	Nonsense	From parents	Pathogenic	Novel
	<i>EIF2AK3</i>	c.1912C>T (p.Arg638*)	Nonsense			
432	<i>SLC19A2</i>	c.164delC (p.Pro55fs)	Frameshift	Mother	Pathogenic	Novel
	<i>SLC19A2</i>	c.161C>A (p.Thr54Asn)	Missense	Father	Uncertain significance	Novel
226	<i>GCK</i>	c.543_545delCGT (p.Val182del)	In-frame deletion	Unknown	Uncertain significance	Novel
	<i>HNF1A</i>	c.92G>A (p.Gly31Asp)	Missense	Unknown	Likely pathogenic	(39)
529	<i>BLK</i>	c.939G>C (p.Glu313Asp)	Missense	Unknown	Uncertain significance	Novel
	<i>GCK</i>	c.919C>T (p.Leu307Phe)	Missense	Unknown	Uncertain significance	Novel
662	<i>GCK</i>	c.1019+2T>A	Splicing defect	Unknown	Pathogenic	Novel
	<i>BLK</i>	c.1148G>A (p.Arg383Gln)	Missense	Unknown	Uncertain significance	Novel
	<i>WFS1</i>	c.1957C>T (p.Arg653Cys)	Missense	Unknown	Likely pathogenic	(40)

ACMG, American College of Medical Genetics and Genomics.

the mother, we concluded that only one allele was affected. Moreover, only nonsense mutation c.199G>T seemed to be clinically significant, because the resulting stop-codon

terminates translation before the c.766G>C site. The clinical picture in patient #226, who had genetic variants in *GCK* and *HNF1A*, was more typical for MODY2 than for MODY3:




Figure 1. The spectrum of genetic variants in the *GCK* gene identified in Russian children with non-type 1 diabetes mellitus. Exons and variants are numbered according to the canonical transcript (ENST00000403799.8). Novel variants are highlighted in red. The lower panel indicates the distribution of known pathogenic and likely pathogenic coding variants in *GCK* according to ClinVar (v. 2019-06-18). P, pathogenic; LP, likely pathogenic.

He had mild fasting and postprandial hyperglycemia, had no glucosuria, and was successfully being treated by a diet. Patient #411 had a homozygous *EIF2AK3* nonsense mutation, inherited from consanguineous parents and associated with Wolcott-Rallison syndrome, which, in turn, has been reported to be the most common genetic cause of permanent neonatal diabetes in consanguineous families (46). Patient #432 had two novel genetic variants affecting both *SLC19A2* alleles. Homozygous mutations in *SLC19A2* cause Rogers syndrome: Thiamine-responsive megaloblastic anaemia associated with diabetes mellitus and deafness (47). Among other clinical signs are congenital heart defects, retinal degeneration, ketonuria, dwarfism, and neurological symptomatology (42). Of note, patient #432 had only diabetes mellitus, retinal degeneration, ketonuria, and neurological symptomatology and thus did not manifest a typical clinical picture. Both patients #529 and #662 had typical clinical signs of *GCK*-MODY rather than *BLK*-MODY, suggesting an absence of strong accumulation of the pathogenic effect of the detected genetic variants.

## Discussion

In 1974, Tattersall reported on three families suffering from mild non-insulin dependent diabetes with Mendelian dominant inheritance (48). The disease was diagnosed in children and young adults and was later defined as maturity-onset type diabetes of young people (MODY) (49). The discovery of mutations in the genes encoding HNF4A (50), HNF1A (51), HNF1B (52), IPF (PDX1) (53), and *GCK* (54,55) as the causes of MODY provided evidence for genetic heterogeneity of familial diabetes. To date, MODY-causing mutations are identified in a total of 13 genes, and mutations in more than 20 genes are known to be associated with neonatal hyperglycemia (56). Because of such a variety of genetic causes, many cases of monogenic diabetes remain without a genetic diagnosis, and its frequency remains underestimated.

The development of high throughput sequencing became a milestone in the search for diabetes-related mutations.

Allowing for simultaneous testing of an unlimited number of genes (i.e. of all known genetic etiology in monogenic diabetes), the method increased the mutation detection rate significantly (57). In our study, we detected genetic variants causative of monogenic diabetes and hyperglycemia-related syndromes in 33 out of 60 children (55%) with non-type 1 diabetes mellitus. This frequency is considerably higher than that detected by Sanger sequencing, which is usually restricted to the analysis of several MODY-related genes and confirms approximately 15% of the cases tested for MODY (58). The higher mutation detection rate in our study is achieved by increasing the number of genes tested and a thorough clinical selection of patients with possible monogenic diabetes. In this regard, one more advantage of WES should be mentioned: DNA sequencing data may be easily stored for further analysis of newly discovered candidate genes.

Ethnic differences play an important role in determining the epidemiology of monogenic diabetes, especially of MODY. Large population studies in European Caucasians showed a general trend of increased *HNF1A*-MODY frequency in Northern Europe, while *GCK*-MODY is prevalent in Southern European populations (5). Here, we report *GCK*-MODY in 19 and *HNF1A*-MODY in only 2 out of 27 MODY-positive Russian patients. These mutation rates appeared to be closer to those in Southern European populations than to those in Northern Europe residents. Our finding may indicate the population-specific frequency MODY types in Russian patients. The recently shown high prevalence of *GCK*-MODY cases among Russian patients with diabetes in pregnancy supports this suggestion (59). However, it should be also considered that our study was performed on children who developed diabetes before the age of 18 years. In the previous observations, it was noticed that the relative proportion of *GCK*-MODY is higher in cases ascertained through pediatric clinics, in contrast to *HNF1A*-MODY, which predominates in cases from adult clinics (58,60). Thus, considering this information, our results are in good accordance with those reported in Spain, Italy, France, Germany, and the Czech Republic, where mostly

Table II. *In silico* prediction of increase/decrease in the protein stability caused by missense mutations with uncertain significance and by benign missense mutations.

Patient number	Gene	Genetic variant (amino acid change)	Pathogenicity according to ACMG	Protein stability predicted by I-Mutant
434	<i>GCK</i>	c.868G>C (p.Glu290Gln)	Uncertain significance	Decrease
153	<i>HNF1A</i>	c.709A>G (p.Asn237Asp)	Uncertain significance	Decrease
422	<i>HNF1A</i>	c.485T>G (p.Leu162Arg)	Uncertain significance	Decrease
215	<i>PAX4</i>	c.574C>A (p.Arg192Ser)	Uncertain significance	Decrease
134	<i>KCNJ11</i>	c.406C>A (p.Arg136Ser)	Uncertain significance	Decrease
266	<i>WFS1</i>	c.2452C>T (p.Arg818Cys)	Likely benign	Decrease
408	<i>WFS1</i>	c.2327A>T (p.Glu776Val)	Likely benign	Increase
133	<i>WFS1</i>	c.1124G>A (p.Arg375His)	Uncertain significance	Decrease
432	<i>SLC19A2</i>	c.161C>A (p.Thr54Asn)	Uncertain significance	Decrease
529	<i>BLK</i>	c.939G>C (p.Glu313Asp)	Uncertain significance	Decrease
	<i>GCK</i>	c.919C>T (p.Leu307Phe)	Uncertain significance	Decrease
662	<i>BLK</i>	c.1148G>A (p.Arg383Gln)	Uncertain significance	Decrease

ASMG, American College of Medical Genetics and Genomics.

pediatric cases were tested (25,61). The prevalence of *GCK* variants (57.6%) in our study suggests that genetic analysis in Russian children with suspected monogenic diabetes may start with testing for *MODY2*, which may not necessarily be performed by WES. However, other cases amount to 42.4% and are linked to 9 different genes, which attests to the efficiency of using WES for the search of genetic causes of diabetes in non-*GCK-MODY* cases.

Our results show that the spectrum of monogenic diabetes-related genetic variants in Russian children includes missense and nonsense mutations, in-frame and frameshift deletions, and a promoter mutation. Generally, these data do not contrast with results obtained in other populations, which also demonstrated a wide spectrum of mutations (62-64). Among genetic variants detected in our study, 60.5% had already been reported in diabetic patients and 39.5% were novel ones. On the one hand, these results point towards a significant recurrent variation within monogenic-diabetes-related genes. On the other hand, they suggest that, in spite of the multitude of monogenic diabetes studies, many variants still remain unidentified. Identification of novel genetic variants as well as accumulating data on previously known causes of monogenic diabetes is of high importance, both for fundamental understanding of the disease pathogenesis and for clinical practice.

Interpretation of genetic variants, especially novel ones, may be challenging. In this study, only 63.2% of the detected genetic variants (18 previously reported and 6 novel ones) were unambiguously considered as causative of the diabetes in the studied patients. The remaining 36.8% variants, including 9 novel ones, were initially classified as those of uncertain significance (n=12) or likely benign (n=2). Additional *in silico* predictions performed for missense mutations among these variants indicate that, with the exception of one variant, they all likely have an adverse effect on protein stability. Considering these results and the patients' phenotypes, the

assumption that the abovementioned variants may be causative of monogenic diabetes can be made. Importantly, the detected genetic variants are absent in non-diabetic Russian population resided in Northwest Russia (65). However, to make a strong conclusion on the pathogenic effect of each novel variant, more data are required, including functional characterization and reports of a specific genetic variant in multiple patients with similar phenotypes. The latter highlights the importance of our results for future studies of monogenic diabetes-related genetic variants.

Noteworthy, our analysis of the clinical picture in the patients simultaneously having *BLK+GCK* (patient #529 and #662) and *GCK+HNF1A* (patient #226) genetic variants suggests no accumulation of adverse effect: All these patients had a typical *MODY2* phenotype. The most plausible explanation for this is the specific age of development of different *MODY* types. Patients suffering from *GCK-MODY* have an impaired glucose metabolism since birth (66). In contrast, carriers of *HNF1A* genetic variants may develop diabetes by the age of 35 years or even by the age of 55 years, although most of them have diabetes before 25 years of age (67). In the study by López-Garrido *et al* (68), the co-inheritance of *GCK* and *HNF1A* genetic variants was reported in two patients and was associated with a typical *MODY3* phenotype in an adult patient and only impaired fasting glucose in a younger patient with the same genotype. In addition, *HNF1A* genetic variant detected in patient #226 in this study (c.92G>A) was previously reported in a diabetic proband and his non-diabetic sister of 43 years of age (69). Similarly, affected carriers of *BLK* genetic variants usually develop diabetes at the middle age (70). Thus, it is likely that patients #529, #662, and #226, who were all involved in our study before the age of 4 years, have not developed the clinical picture of *HNF1A-MODY* and *BLK-MODY* yet. The possibility of a late manifestation of *HNF1A-MODY* and *BLK-MODY* in the children who

Table III. Clinical characteristics of the patients with multiple genetic variants in monogenic diabetes-related genes.

Patient number	Gene Nucleotide change Amino acid change	Age at diagnosis months	Diabetic ketoacidosis	C-peptide ng/ml	HbA1C %	SDS BMI	Treatment
27	<i>GCK</i> c.754T>C (p.Cys252Arg) <i>GCK</i> c.-71G>C	3	No	0.7	6	-0.63	Diet
78	<i>GCK</i> c.199G>T (p.Glu67*) <i>GCK</i> c.766G>C (p.Glu256Lys)	39	No	0.63	6.4	+0.83	Diet
226	<i>GCK</i> c.543_545delCGT (p.Val182del) <i>HNF1A</i> c.92G>A (p.Gly31Asp)	36	No	1.1	6	-1.69	Diet
411	<i>EIF2AK3</i> c.1912C>T (p.Arg638*) <i>EIF2AK3</i> c.1912C>T (p.Arg638*)	3	Ketonuria	0.2	9.2	-0.72	Insulin
432	<i>SLC19A2</i> c.164delC (p.Pro55fs) <i>SLC19A2</i> c.161C>A (p.Thr54Asn)	48	Ketonuria	1.1	5.3	-1.0	Insulin for a few days/ diet
529	<i>BLK</i> c.939G>C (p.Glu313Asp) <i>GCK</i> c.919C>T (p.Leu307Phe)	10	No	0.43	6.7	-0.46	Diet
662	<i>GCK</i> c.1019+2T>A <i>BLK</i> c.1148G>A (p.Arg383Gln) <i>WFS1</i> c.1957C>T (p.Arg653Cys)	22	No	1.1	6.82	-1.32	Diet

SDS BMI reference range: -1.5/+1.5; SDS, standard deviation score.

already have *GCK*-MODY strongly suggests the necessity of their strict medical supervision in order to timely modify their therapy. Additional studies, including functional ones, on the pathogenicity of the novel *BLK* genetic variants detected in patients #529 and #662 will also facilitate the development of the most effective treatment strategies for them.

To summarize, our data show a high rate of genetic variants causative of monogenic diabetes in Russian children with non-type 1 diabetes mellitus. The use of a WES-based panel allowed us to identify a variety of previously known and novel genetic variants in MODY-related and unrelated genes, including multiple variants in a number of patients. The revealed variety is characterized by a prevalence of *GCK* genetic variants (MODY2) and also includes variants in *HNF1A*, *PAX4*, *KCNJ11*, *BLK*, *ABCC8*, *GATA6*, *WFS1*, *EIF2AK3*, and *SLC19A2*. These results, on the one hand, suggest that genetic analysis for monogenic diabetes in Russian children may start

with testing for *GCK* variants, which may not necessarily be performed by WES. On the other hand, non-*GCK* variants are linked to 9 different genes, which attests to the efficiency of using WES while searching for genetic causes of diabetes in non-*GCK*-MODY cases. Notably, the detection of genetic variants in the genes linked to specific syndromes with recessive inheritance-*WFS1*, *EIF2AK3*, and *SLC19A2*-is essential for appropriate genetic counseling and family planning. Our study highlights the importance of using WES for monogenic diabetes testing and provides new information on the diabetes-related genetic variants in the Russian population.

#### Acknowledgements

The authors would like to thank Mrs Ksenia Khudadyan (Logrus LLC) for helpful advice during the preparation of the manuscript.

## Funding

The current study was supported by the Ministry of Science and Higher Education of the Russian Federation within the framework of the Basic Research Program in 2019-2021 (grant no. AAAA-A19-119021290033-1), the alpha-Endo program, the Charities Aid Foundation Foundation (grant nos. 65/315 and 133/315), and the Russian Science Foundation (grant no. 14-50-00069).

## Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Authors' contributions

OSG, EAS, MET, OAE, EBB and VSB designed the study. OSG, EAS, MET, ASG, YAN, DEP, MAF, IVP, TEI, NYS, ESS, AVT, OVR, AMS, AAP, SGS, EVM, AVPK, LRL, LVD, LAZ, LVT, OSB, ENS and EBB recruited the patients and performed experimental procedures. YAB, AVP and RKS performed bioinformatic analysis. OSG, EAS, MET, OAE, AAP, TEI, LRL, EBB and VSB analyzed result and performed literature search. OAE wrote the manuscript. All authors read and approved the final manuscript.

## Ethics approval and consent to participate

The study was approved by the Ethics Committee of D.O. Ott Research Institute of Obstetrics, Gynecology and Reproductology. All the patients/patients' representatives gave written informed consent to participate in the study. The study was performed in accordance with the Declaration of Helsinki.

## Patient consent for publication

All the patients/patients' representatives gave written informed consent for publication of the study results.

## Competing interests

The authors declare that they have no competing interests.

## References

- Hattersley AT, Greeley SAW, Polak M, Rubio-Cabezas O, Njølstad PR, Mlynarski W, Castano L, Carlsson A, Raile K, Chi DV, *et al*: ISPAD clinical practice consensus guidelines 2018: The diagnosis and management of monogenic diabetes in children and adolescents. *Pediatr Diabetes* 19 (Suppl 27): S47-S63, 2018.
- Barbetti F and D'Annunzio G: Genetic causes and treatment of neonatal diabetes and early childhood diabetes. *Best Pract Res Clin Endocrinol Metab* 32: 575-591, 2018.
- Lemelman MB, Letourneau L and Greeley SAW: Neonatal diabetes mellitus: An update on diagnosis and management. *Clin Perinatol* 45: 41-59, 2018.
- Greeley SA, Naylor RN, Philipson LH and Bell GI: Neonatal diabetes: An expanding list of genes allows for improved diagnosis and treatment. *Curr Diab Rep* 11: 519-532, 2011.
- Kleinberger JW and Pollin TI: Undiagnosed MODY: Time for Action. *Curr Diab Rep* 15: 110, 2015.
- American Diabetes Association: 2. Classification and diagnosis of diabetes: Standards of medical care in diabetes-2018. *Diabetes Care* 41 (Suppl 1): S13-S27, 2018.
- DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del Angel G, Rivas MA, Hanna M, *et al*: A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet* 43: 491-498, 2011.
- Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, Jordan T, Shakir K, Roazen D, Thibault J, *et al*: From FastQ data to high-confidence variant calls: The genome analysis toolkit best practices pipeline. *Curr Protoc Bioinform* 43: 11.10.1-33, 2013.
- 1000 Genomes Project Consortium; Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA and Abecasis GR: A global reference for human genetic variation. *Nature* 526: 68-74, 2015.
- Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, O'Donnell-Luria AH, Ware JS, Hill AJ, Cummings BB, *et al*: Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 536: 285-291, 2016.
- Fu W, O'Connor TD, Jun G, Kang HM, Abecasis G, Leal SM, Gabriel S, Rieder MJ, Altshuler D, Shendure J, *et al*: Analysis of 6,515 exomes reveals the recent origin of most human protein-coding variants. *Nature* 493: 216-220, 2013.
- Adzhubei IA, Schmidt S, Peshkin S, Ramensky VE, Gerasimova A, Bork P, Kondrashov AS and Sunyaev SR: A method and server for predicting damaging missense mutations. *Nat Methods* 7: 248-249, 2010.
- Ng PC and Henikoff S: Predicting deleterious amino acid substitutions. *Genome Res* 11: 863-874, 2001.
- Choi Y and Chan AP: PROVEAN web server: A tool to predict the functional effect of amino acid substitutions and indels. *Bioinformatics* 31: 2745-2747, 2015.
- Liu X, Jian X and Boerwinkle E: dbNSFP v2.0: A database of human non-synonymous SNVs and their functional predictions and annotations. *Hum Mutat* 34: E2393-E2402, 2013.
- Desmet FO, Hamroun D, Lalonde M, Collod-Bérout G, Claustres M and Bérout C: Human splicing finder: An online bioinformatics tool to predict splicing signals. *Nucleic Acids Res* 37: e67, 2009.
- Zhao H, Yang Y, Lin H, Zhang X, Mort M, Cooper DN, Liu Y and Zhou Y: DDIG-in: Discriminating between disease-associated and neutral non-frameshifting micro-indels. *Genome Biol* 14: R23, 2013.
- Capriotti E, Fariselli P and Casadio R: I-Mutant2.0: Predicting stability changes upon mutation from the protein sequence or structure. *Nucleic Acids Res* 33: W306-W310, 2005.
- Barbitoff YA, Bezdvornykh IV, Polev DE, Serebryakova EA, Glotov AS, Glotov OS and Predeus AV: Catching hidden variation: Systematic correction of reference minor allele annotation in clinical variant calling. *Genet Med* 20: 360-364, 2018.
- Gasperičková D, Tribble ND, Staník J, Hucková M, Misovicová N, van de Bunt M, Valentínová L, Barrow BA, Barák L, Dobránský R, *et al*: Identification of a novel beta-cell glucokinase (GCK) promoter mutation (-71G>C) that modulates GCK gene expression through loss of allele-specific Sp1 binding causing mild fasting hyperglycemia in humans. *Diabetes* 58: 1929-1935, 2009.
- Mantovani V, Salardi S, Cerreta V, Bastia D, Cenci M, Ragni L, Zucchini S, Parente R and Cicognani A: Identification of eight novel glucokinase mutations in Italian children with maturity-onset diabetes of the young. *Hum Mutat* 22: 338, 2003.
- Bacon S, Kyithar MP, Schmid J, Rizvi SR, Bonner C, Graf R, Prehn JH and Byrne MM: Serum levels of pancreatic stone protein (PSP)/reg1A as an indicator of beta-cell apoptosis suggest an increased apoptosis rate in hepatocyte nuclear factor 1 alpha (HNF1A-MODY) carriers from the third decade of life onward. *BMC Endocr Disord* 12: 13, 2012.
- Graglioli C, Cockburn BN, Chiaramonte F, Gorini A, Marietti G, Marozzi G and Signorini AM: Early-onset Type II diabetes mellitus in Italian families due to mutations in the genes encoding hepatic nuclear factor 1 alpha and glucokinase. *Diabetologia* 44: 1326-1329, 2001.
- Ziemssen F, Bellanné-Chantelot C, Osterhoff M, Schatz H and Pfeiffer AF: To: Lindner T, cockburn BN, Bell GI (1999) Molecular genetics of MODY in Germany. *Diabetologia* 42: 121-123. *Diabetologia* 45: 286-288, 2002.
- Estalella I, Rica I, Perez de Nancrales G, Bilbao JR, Vazquez JA, San Pedro JI, Busturia MA and Castaño L: Spanish MODY Group: Mutations in GCK and HNF-1alpha explain the majority of cases with clinical diagnosis of MODY in Spain. *Clin Endocrinol (Oxf)* 67: 538-546, 2007.

26. Stoffel M, Froguel P, Takeda J, Zouali H, Vionnet N, Nishi S, Weber IT, Harrison RW, Pilkis SJ, Lesage S, *et al*: Human glucokinase gene: Isolation, characterization, and identification of two missense mutations linked to early-onset non-insulin-dependent (type 2) diabetes mellitus. *Proc Natl Acad Sci USA* 89: 7698-7702, 1992.
27. Hattersley AT, Beards F, Ballantyne E, Appleton M, Harvey R and Ellard S: Mutations in the glucokinase gene of the fetus result in reduced birth weight. *Nat Genet* 19: 268-270, 1998.
28. Froguel P, Zouali H, Vionnet N, Velho G, Vaxillaire M, Sun F, Lesage S, Stoffel M, Takeda J, Passa P, *et al*: Familial hyperglycemia due to mutations in glucokinase. Definition of a subtype of diabetes mellitus. *N Engl J Med* 328: 697-702, 1993.
29. Borowiec M, Fendler W, Antosik K, Baranowska A, Gnys P, Zmyslowska A, Malecki M and Mlynarski W: Doubling the referral rate of monogenic diabetes through a nationwide information campaign-update on glucokinase gene mutations in a Polish cohort. *Clin Genet* 82: 587-590, 2012.
30. Pruhova S, Ek J, Lebl J, Sumnik Z, Saudek F, Andel M, Pedersen O and Hansen T: Genetic epidemiology of MODY in the Czech republic: New mutations in the MODY genes HNF-4alpha, GCK and HNF-1alpha. *Diabetologia* 46: 291-295, 2003.
31. Gidh-Jain M, Takeda J, Xu LZ, Lange AJ, Vionnet N, Stoffel M, Froguel P, Velho G, Sun F, Cohen D, *et al*: Glucokinase mutations associated with non-insulin-dependent (type 2) diabetes mellitus have decreased enzymatic activity: Implications for structure/function relationships. *Proc Natl Acad Sci USA* 90: 1932-1936, 1993.
32. Colclough K, Bellanne-Chantelot C, Saint-Martin C, Flanagan SE and Ellard S: Mutations in the genes encoding the transcription factors hepatocyte nuclear factor 1 alpha and 4 alpha in maturity-onset diabetes of the young and hyperinsulinemic hypoglycemia. *Hum Mutat* 34: 669-685, 2013.
33. Plengvidhya N, Kooptiwut S, Songtawe N, Doi A, Furuta H, Nishi M, Nanjo K, Tantibhedhyangkul W, Boonyasrisawat W, Yenchitsomanus PT, *et al*: PAX4 mutations in Thais with maturity onset diabetes of the young. *J Clin Endocrinol Metab* 92: 2821-2826, 2007.
34. Flanagan SE, Patch AM, Mackay DJ, Edghill EL, Gloyn AL, Robinson D, Shield JP, Temple K, Ellard S and Hattersley AT: Mutations in ATP-sensitive K<sup>+</sup> channel genes cause transient neonatal diabetes and permanent diabetes in childhood or adulthood. *Diabetes* 56: 1930-1937, 2007.
35. Mohnike K, Wieland I, Barthlen W, Vogelgesang S, Empting S, Mohnike W, Meissner T and Zenker M: Clinical and genetic evaluation of patients with KATP channel mutations from the German registry for congenital hyperinsulinism. *Horm Res Paediatr* 81: 156-168, 2014.
36. Suzuki S, Nakao A, Sarhat AR, Furuya A, Matsuo K, Tanahashi Y, Kajino H and Azuma H: A case of pancreatic agenesis and congenital heart defects with a novel GATA6 nonsense mutation: Evidence of haploinsufficiency due to nonsense-mediated mRNA decay. *Am J Med Genet A* 164A: 476-479, 2014.
37. Gomez-Zaera M, Strom T, Meitinger T and Nunes V: Wolframin mutations in Spanish families with Wolfram syndrome. Meeting abstract. *Am J Hum Genet* 65: 1673, 1999.
38. Smith CJ, Crock PA, King BR, Meldrum CJ and Scott RJ: Phenotype-genotype correlations in a series of wolfram syndrome families. *Diabetes Care* 27: 2003-2009, 2004.
39. Chèvre JC, Hani EH, Boutin P, Vaxillaire M, Blanché H, Vionnet N, Pardini VC, Timsit J, Larger E, Charpentier G, *et al*: Mutation screening in 18 Caucasian families suggest the existence of other MODY genes. *Diabetologia* 41: 1017-1023, 1998.
40. Awata T, Inoue K, Kurihara S, Ohkubo T, Inoue I, Abe T, Takino H, Kanazawa Y and Katayama S: Missense variations of the gene responsible for Wolfram syndrome (WFS1/wolframin) in Japanese: Possible contribution of the Arg456His mutation to type 1 diabetes as a nonautoimmune genetic basis. *Biochem Biophys Res Commun* 268: 612-616, 2000.
41. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M, Lyon E, Spector E, *et al*: Standards and guidelines for the interpretation of sequence variants: A joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med* 17: 405-424, 2015.
42. Shaw-Smith C, Flanagan SE, Patch AM, Grulich-Henn J, Habeb AM, Hussain K, Pomahacova R, Matyka K, Abdullah M, Hattersley AT and Ellard S: Recessive SLC19A2 mutations are a cause of neonatal diabetes mellitus in thiamine-responsive megaloblastic anaemia. *Pediatr Diabetes* 13: 314-321, 2012.
43. Fraser FC and Gunn T: Diabetes mellitus, diabetes insipidus, and optic atrophy. An autosomal recessive syndrome? *J Med Genet* 14: 190-193, 1977.
44. Bennett K, James C, Mutair A, Al-Shaikh H, Sinani A and Hussain K: Four novel cases of permanent neonatal diabetes mellitus caused by homozygous mutations in the glucokinase gene. *Pediatr Diabetes* 12: P192-P196, 2011.
45. Raimondo A, Chakera AJ, Thomsen SK, Colclough K, Barrett A, De Franco E, Chatelas A, Demirbilek H, Akcay T, Alawneh H, *et al*: Phenotypic severity of homozygous GCK mutations causing neonatal or childhood-onset diabetes is primarily mediated through effects on protein stability. *Hum Mol Genet* 23: 6432-6440, 2014.
46. Rubio-Cabezas O, Patch AM, Minton JA, Flanagan SE, Edghill EL, Hussain K, Balafrej A, Deeb A, Buchanan CR, Jefferson IG, *et al*: Wolcott-Rallison syndrome is the most common genetic cause of permanent neonatal diabetes in consanguineous families. *J Clin Endocrinol Metab* 94: 4162-4170, 2009.
47. Labay V, Raz T, Baron D, Mandel H, Williams H, Barrett T, Szargel R, McDonald L, Shalata A, Nosaka K, *et al*: Mutations in SLC19A2 cause thiamine-responsive megaloblastic anaemia associated with diabetes mellitus and deafness. *Nat Genet* 22: 300-304, 1999.
48. Tattersall RB: Mild familial diabetes with dominant inheritance. *Q J Med* 43: 339-357, 1974.
49. Tattersall RB and Fajans SS: A difference between the inheritance of classical juvenile-onset and maturity-onset type diabetes of young people. *Diabetes* 24: 44-53, 1975.
50. Yamagata K, Furuta H, Oda N, Kaisaki PJ, Menzel S, Cox NJ, Fajans SS, Signorini S, Stoffel M and Bell GI: Mutations in the hepatocyte nuclear factor-4alpha gene in maturity-onset diabetes of the young (MODY1). *Nature* 384: 458-460, 1996.
51. Yamagata K, Oda N, Kaisaki PJ, Menzel S, Furuta H, Vaxillaire M, Southam L, Cox RD, Lathrop GM, Boriraj VV, *et al*: Mutations in the hepatocyte nuclear factor-1alpha gene in maturity-onset diabetes of the young (MODY3). *Nature* 384: 455-458, 1996.
52. Horikawa Y, Iwasaki N, Hara M, Furuta H, Hinokio Y, Cockburn BN, Lindner T, Yamagata K, Ogata M, Tomonaga O, *et al*: Mutation in hepatocyte nuclear factor-1 beta gene (TCF2) associated with MODY. *Nat Genet* 17: 384-385, 1997.
53. Stoffers DA, Ferrer J, Clarke WL and Habener JF: Early-onset type-II diabetes mellitus (MODY4) linked to IPF1. *Nat Genet* 17: 138-139, 1997.
54. Froguel P, Vaxillaire M, Sun F, Velho G, Zouali H, Butel MO, Lesage S, Vionnet N, Clément K, Fougerousse F, *et al*: Close linkage of glucokinase locus on chromosome 7p to early-onset non-insulin-dependent diabetes mellitus. *Nature* 356: 162-164, 1992.
55. Hattersley AT, Turner RC, Permutt MA, Patel P, Tanizawa Y, Chiu KC, O'Rahilly S, Watkins PJ and Wainscoat JS: Linkage of type 2 diabetes to the glucokinase gene. *Lancet* 339: 1307-1310, 1992.
56. Timsit J, Saint-Martin C, Dubois-Laforgue D and Bellanné-Chantelot C: Searching for Maturity-Onset diabetes of the Young (MODY): When and What for? *Can J Diabetes* 40: 455-461, 2016.
57. Ellard S, Lango Allen H, De Franco E, Flanagan SE, Hysenaj G, Colclough K, Houghton JA, Shepherd M, Hattersley AT, Weedon MN and Caswell R: Improved genetic testing for monogenic diabetes using targeted next-generation sequencing. *Diabetologia* 56: 1958-1963, 2013.
58. Shields BM, Hicks S, Shepherd MH, Colclough K, Hattersley AT and Ellard S: Maturity-onset diabetes of the young (MODY): How many cases are we missing? *Diabetologia* 53: 2504-2508, 2010.
59. Zubkova N, Burumkulova F, Plechanova M, Petrukhin V, Petrov V, Vasilyev E, Panov A, Sorkina E, Ulyatovskaya V, Makretskaya N and Tiulpakov A: High frequency of pathogenic and rare sequence variants in diabetes-related genes among Russian patients with diabetes in pregnancy. *Acta Diabetol* 56: 413-420, 2019.
60. Lorini R, Klersy C, d'Annunzio G, Massa O, Minuto N, Iafusco D, Bellanné-Chantelot C, Frongia AP, Toni S, Meschi F, *et al*: Maturity-onset diabetes of the young in children with incidental hyperglycemia: A multicenter Italian study of 172 families. *Diabetes Care* 32: 1864-1866, 2009.
61. Schober E, Rami B, Grabert M, Thon A, Kapellen T, Reinehr T and Holl RW: Phenotypical aspects of maturity-onset diabetes of the young (MODY diabetes) in comparison with Type 2 diabetes mellitus (T2DM) in children and adolescents: Experience from a large multicentre database. *Diabet Med* 26: 466-473, 2009.

62. Ağladioğlu SY, Aycan Z, Çetinkaya S, Baş VN, Önder A, Peltek Kendirci HN, Doğan H and Ceylaner S: Maturity onset diabetes of youth (MODY) in Turkish children: Sequence analysis of 11 causative genes by next generation sequencing. *J Pediatr Endocrinol Metab* 29: 487-496, 2016.
63. Globa E, Zelinska N, Elblova L, Dusatkova P, Cinek O, Lebl J, Colclough K, Ellard S and Pruhova S: MODY in Ukraine: Genes, clinical phenotypes and treatment. *J Pediatr Endocrinol Metab* 30: 1095-1103, 2017.
64. Johansson BB, Irgens HU, Molnes J, Sztromwasser P, Aukrust I, Juliusson PB, Søvik O, Levy S, Skriverhaug T, Joner G, *et al*: Targeted next-generation sequencing reveals MODY in up to 6.5% of antibody-negative diabetes cases listed in the Norwegian Childhood Diabetes Registry. *Diabetologia* 60: 625-635, 2017.
65. Barbitoff YA, Skitchenko RK, Poleshchuk OI, Shikov AE, Serebryakova EA, Nasykhova YA, Polev DE, Shuvalova AR, Shcherbakova IV, Fedyakov MA, *et al*: Whole-exome sequencing provides insights into monogenic disease prevalence in Northwest Russia. *Mol Genet Genomic Med*: Sep 3: e964, 2019 doi: 10.1002/mgg3.964 (Epub ahead of print).
66. Chakera AJ, Steele AM, Gloyn AL, Shepherd MH, Shields B, Ellard S and Hattersley AT: Recognition and management of individuals with hyperglycemia because of a heterozygous glucokinase mutation. *Diabetes Care* 38: 1383-1392, 2015.
67. Murphy R, Ellard S and Hattersley AT: Clinical implications of a molecular genetic classification of monogenic beta-cell diabetes. *Nat Clin Pract Endocrinol Metab* 4: 200-213, 2008.
68. López-Garrido MP, Herranz-Antolín S, Alija-Merillas MJ, Giralt P and Escribano J: Co-inheritance of HNF1a and GCK mutations in a family with maturity-onset diabetes of the young (MODY): Implications for genetic testing. *Clin Endocrinol (Oxf)* 79: 342-347, 2013.
69. Thanabalasingham G, Pal A, Selwood MP, Dudley C, Fisher K, Bingley PJ, Ellard S, Farmer AJ, McCarthy MI and Owen KR: Systematic assessment of etiology in adults with a clinical diagnosis of young-onset type 2 diabetes is a successful strategy for identifying maturity-onset diabetes of the young. *Diabetes Care* 35: 1206-1212, 2012.
70. Borowiec M, Liew CW, Thompson R, Boonyasrisawat W, Hu J, Mlynarski WM, El Khattabi I, Kim SH, Marselli L, Rich SS, *et al*: Mutations at the BLK locus linked to maturity onset diabetes of the young and beta-cell dysfunction. *Proc Natl Acad Sci USA* 106: 14460-14465, 2009.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.